

# Quadric surfaces fibrations over the real projective line joint work with Alena Pirutka

Jean-Louis Colliot-Thélène  
CNRS et Université Paris-Saclay

Workshop on algebraic cycles  
30. September – 2. Oktober 2024  
Leibniz Universität Hannover

Basic rational varieties over an algebraically closed field : quadrics ; total space of a family of positive dimensional quadrics over the projective line.

A basic birational invariant of smooth, projective, geometrically connected varieties  $X$  over the field  $\mathbb{R}$  : the number of connected components of the topological space  $X(\mathbb{R})$ .

Beniamino Segre (1951) : a smooth real cubic surface  $X \subset \mathbb{P}_{\mathbb{R}}^3$  is  $\mathbb{R}$ -unirational, is  $\mathbb{C}$ -rational, but if  $X(\mathbb{R})$  has two connected components, it is not  $\mathbb{R}$ -rational.

A smooth, projective, geometrically rational surface  $X$  over  $\mathbb{R}$  is  $\mathbb{R}$ -rational if and only if  $X(\mathbb{R})$  is nonempty and connected (Comessatti 1913, Silhol 1989). Proof uses birational classification.

Suppose  $X(\mathbb{R})$  nonempty and connected and  $X$  is rational over  $\mathbb{C}$ .

Is  $X$   $\mathbb{R}$ -rational?

(Weaker)

Is  $X$  stably  $\mathbb{R}$ -rational?

(Weaker)

Is  $X$  universally Chow-trivial : for any overfield  $F/\mathbb{R}$ , is the degree map  $deg_F : CH_0(X_F) \rightarrow \mathbb{Z}$  an isomorphism? (Equivalently, there is a decomposition of the diagonal for  $X/\mathbb{R}$ )

Enough to prove it for  $F = \mathbb{R}(X)$ .

Over any field  $k$ , the question of rationality over  $k$  of geometrically rational threefolds  $X/k$  has been the topic of works of Hassett-Tschinkel and Benoist-Wittenberg. The IJT (intermediate Jacobian torsor) method is an extension to arbitrary ground fields of the method used by Clemens-Griffiths to disprove rationality of cubic threefolds over  $\mathbb{C}$ .

Let us consider  $X = X_{2,2} \subset \mathbb{P}_{\mathbb{R}}^5$  a smooth complete intersection of two quadrics. Over  $\mathbb{C}$ , rational. Over  $\mathbb{R}$ , the IJT method gives non- $\mathbb{R}$ -rationality of  $X$  when  $X$  contains no real line (may happen with  $X(\mathbb{R})$  connected). It does not give information on stable rationality or universal  $CH_0$ -triviality.

Let  $X \rightarrow \mathbb{P}_{\mathbb{R}}^1$  be a family of quadrics of relative dimension  $d \geq 1$ .  
with smooth total space  $X/\mathbb{R}$ .

$X_{\mathbb{C}}$  is rational. If  $X(\mathbb{R})$  is nonempty,  $X$  is  $\mathbb{R}$ -unirational (exercise).

If  $X(\mathbb{R})$  is connected, is  $X$  rational over  $\mathbb{R}$ ? Is it at least stably rational? Is it universally  $CH_0$ -trivial?

For  $d = 1$ , yes for all questions. We shall consider the case  $d = 2$ ,  
quadric surface fibrations.

*First negative answer,  $d = 2$*

Let  $X$  be a smooth projective model of the variety given by the equation

$$x^2 + (1 + u^2)y^2 - u(z^2 + t^2) = 0$$

in  $\mathbb{P}^3 \times \mathbb{A}^1$ , coordinates  $(x, y, z, t; u)$ .

Real points cover exactly  $u \geq 0$ , real fibres are connected. This gives connectedness of the real locus. Over  $u = 0$  and over  $u = \infty$  the family degenerates to two conjugate planes. One computes  $\text{Br}(X)/\text{Br}(\mathbb{R}) \neq 0$ . The class  $(-1, u) \in \text{Br}(\mathbb{R}(X))$  is not in the image of  $\text{Br}(\mathbb{R})$  and is unramified.

This implies that  $X$  is not universally  $CH_0$ -trivial, hence not  $\mathbb{R}$ -rational and not even stably  $\mathbb{R}$ -rational.

Definition (for this talk).

Let  $k$  be a field of char. 0. Let  $X \rightarrow \mathbb{P}_k^1$  be a quadric surface fibration. We call it a **good fibration** if  $X/k$  is smooth projective and all the geometric fibres of  $X \rightarrow \mathbb{P}_k^1$  are geometrically integral.

In other words, the fibres are smooth or cones over a smooth conic.

The previous example is not a good fibration.

For good fibrations,  $\text{Br}(k) \rightarrow \text{Br}(X)$  is onto.

## A reminder on quadric surface fibrations

Given a field  $k$ , say of char. zero, and a quadric surface fibration  $X \rightarrow \mathbb{P}_k^1$  the generic fibre is a smooth quadric  $Y \subset \mathbb{P}_K^3$  where  $K = k(\mathbb{P}^1)$ . Let  $q/K$  be a 4-dimensional quadratic form defining  $Y$ . Let  $L/K$  be the quadratic extension defined by the discriminant of  $q$ .

Thus  $L = k(\Delta)$  for  $\Delta$  a smooth projective curve over  $k$ . For any smooth plane (conic) section  $Z/K$  of  $Y \subset \mathbb{P}_K^3$ , the associated quaternion class in  $\text{Br}(K)$  has image a class  $\beta \in \text{Br}(L)$  uniquely defined by  $X/\mathbb{P}_k^1$ . We have  $Y \simeq R_{L/K}(Z_L)$ .

If  $X/\mathbb{P}_k^1$  is a good fibration, we have  $\beta \in \text{Br}(\Delta)$ .



*Back to the problem : Second negative answer,  $d = 2$*

What about good fibrations ?

In the 2021 paper by Hassett and Tschinkel, one finds examples of smooth  $Y_{2,2} \subset \mathbb{P}_{\mathbb{R}}^5$  with  $Y(\mathbb{R})$  connected and  $Y$  not  $\mathbb{R}$ -rational because of IJT obstruction.

One may birationally transform this into a good quadric surface fibration  $X \rightarrow \mathbb{P}_{\mathbb{R}}^1$  with 6 geometric singular fibres. Here the quotient  $\text{Br}(X)/\text{Br}(\mathbb{R}) = 0$ ,  $X(\mathbb{R})$  is connected and  $X$  is not rational over  $\mathbb{R}$ .

Under the **additional assumption** :

*The class of  $\beta \in \text{Br}(\Delta) \subset \text{Br}(k(\Delta))$  is not in the image of  $\text{Br}(k) \rightarrow \text{Br}(k(\Delta))$*

Wittenberg (2023) applies the IJT method to quadric surface fibrations  $X/\mathbb{P}_k^1$  with at least 6 geometric singular fibres, with application to the problem over the reals.

For instance  $X/\mathbb{P}_{\mathbb{R}}^1$  given by the affine equation

$$(u^2 - 1)x^2 + (u^2 - 2)y^2 + (u^2 - 3)z^2 = 1$$

is not rational over  $\mathbb{R}$  but  $X(\mathbb{R})$  is connected.

The IJT method does not work without the additional assumption.

The following problems remain :

Suppose  $p : X \rightarrow \mathbb{P}_{\mathbb{R}}^1$  is a *good fibration* Suppose  $X(\mathbb{R})$  is connected.

- (1) Is  $X/\mathbb{R}$  stably rational? Is it universally  $CH_0$ -trivial?
- (2) If the *additional assumption* fails, is  $X$  rational over  $\mathbb{R}$ ?

In this talk, we shall concentrate on a very concrete case.

Let  $p(u) \in \mathbb{R}[u]$  a monic, nonconstant, even degree, separable, polynomial, strictly positive on  $\mathbb{R}$ . One easily constructs a good family of quadric surfaces  $X/\mathbb{P}_{\mathbb{R}}^1$  which is a birational model of the affine variety with equation

$$x^2 + y^2 + z^2 = u \cdot p(u),$$

with the projection given by the  $u$  coordinate. For the purpose of this talk, we shall call such a fibration a **special quadric fibration**. The space  $X(\mathbb{R})$  is connected. The curve  $\Delta$  is given by  $w^2 = v \cdot p(-v)$ . *The additional assumption fails.* Indeed  $\beta$  is given by  $(-1, -1)$ . So the IJT method may not be used.

Is  $X$  rational over  $\mathbb{R}$ ? is  $X$  stably rational over  $\mathbb{R}$ ? Is it universally  $CH_0$ -trivial?

Theorem Let  $X/\mathbb{P}_{\mathbb{R}}^1$  be a special quadric fibration with affine equation

$$x^2 + y^2 + z^2 = u \cdot p(u).$$

Let  $\Delta/\mathbb{R}$  be the smooth projective curve with affine equation

$$w^2 = v \cdot p(-v).$$

Let  $W/\mathbb{R}$  the fourfold given by  $W := X \times_{\mathbb{R}} \Delta$ .

(1) The cup-product  $(u + v, -1, -1) \in H^3(\mathbb{R}(W), \mathbb{Z}/2)$  is unramified over  $\mathbb{R}$ . It vanishes if and only if the rational function  $u + v \in \mathbb{R}(W)$  (a sum of 6 squares) is a sum of 4 squares in  $\mathbb{R}(W)$ .

(2) The following conditions are equivalent :

(2a) The variety  $X$  is universally  $CH_0$ -trivial.

(2b) For  $F = \mathbb{R}(\Delta)$ , the map  $H^3(F, \mathbb{Z}/2) \rightarrow H_{nr}^3(F(X)/F, \mathbb{Z}/2)$  is an isomorphism.

(2c) The rational function  $u + v \in \mathbb{R}(W)$  is a sum of 4 squares.

Ingredients of the proof.

Results in algebraic K-theory and quadratic forms (Arason, Merkurjev, Suslin, Kahn-Rost-Sujatha).

Study of Chow groups of zero-cycles for threefolds fibred into quadric surfaces over the projective line  
[CTSk] CT-Skorobogatov, J. K-Theory 7 (1993).

Let  $A_0(X) \subset CH_0(X)$  the group of degree zero cycle classes. Let  $X/\mathbb{P}_k^1$  be a good nonconstant quadric surface fibration  $X/\mathbb{P}_k^1$ . Let the double cover  $\Delta/\mathbb{P}_k^1$  and  $\beta \in \text{Br}(\Delta)[2] \subset H^2(k(\Delta), \mathbb{Z}/2)$  as above. .

[CTSk] gives an **injection**

$$\Phi : A_0(X) \hookrightarrow H_{nr}^3(k(\Delta)/k, \mathbb{Z}/2)/[H^1(k, \mathbb{Z}/2) \cup (\beta)].$$

Let  $X \rightarrow \mathbb{P}_{\mathbb{R}}^1$  be a special quadric fibration. Here  $\Delta(\mathbb{R}) \neq \emptyset$  and  $\beta = (-1, -1)_{\mathbb{R}}$  (the *additional assumption* fails). Let  $W = X \times_{\mathbb{R}} \Delta$ . Using precisely this, taking  $k = \mathbb{R}(X)$ , using [CTSk], we prove :  
**the image under  $\Phi$  of the difference between the generic point of  $X$  and an  $\mathbb{R}$ -rational point above  $u = \infty$  vanishes (that is,  $X/\mathbb{R}$  is universally  $CH_0$ -trivial, that is, there is a decomposition of the diagonal for  $X/\mathbb{R}$ ) if and only if  $(u + v, -1, -1) = 0 \in H^3(\mathbb{R}(W), \mathbb{Z}/2)$ , if and only if  $u + v$  is a sum of 4 squares in  $\mathbb{R}(W)$ .**

**Proving universal  $CH_0$ -triviality via (2c) :**  
**hypotheses on  $p(u)$  ensuring that**  
 **$u + v \in \mathbb{R}(W)$  is a sum of 4 squares**



## $CH_0$ -triviality via sums of 4 squares, $p(u)$ of degree 2

Theorem. Let  $p(u) = u^2 + au + b \in \mathbb{R}[u]$  be separable and nonnegative. Let  $X/\mathbb{P}_{\mathbb{R}}^1$  be a special quadric fibration with affine equation

$$x^2 + y^2 + z^2 = u \cdot p(u).$$

If  $b \geq a^2/3$ , then  $X$  is universally  $CH_0$ -trivial.

This covers the case  $p(u) = u \cdot (u^2 + 1)$  but does not cover the range  $a^2/3 > b > a^2/4$ .

Proof. Recall that  $\Delta$  is defined by  $w^2 = v \cdot p(-v)$ , and  $W = X \times_{\mathbb{R}} \Delta$ . In  $\mathbb{R}(u, v)$ ,

$$\begin{aligned} up(u) + vp(-v) &= (u + v)(u^2 - uv + v^2 + au - av + b) = \\ &= (u + v) \left( \left( u + \frac{a - v}{2} \right)^2 + \frac{3}{4} \left( v - \frac{a}{3} \right)^2 + b - \frac{a^2}{3} \right). \end{aligned}$$

Since  $b - \frac{a^2}{3} \geq 0$ ,  $\frac{up(u) + vp(-v)}{u + v}$  is a sum of 3 squares in  $\mathbb{R}(u, v)$ , hence a sum of 4 squares.

In  $\mathbb{R}(W)$ , we have

$$x^2 + y^2 + z^2 + w^2 = up(u) + vp(-v).$$

Thus  $u + v$  is a sum of 4 squares in  $\mathbb{R}(W)$ .

## $CH_0$ -triviality via sums of 4 squares, $p(u)$ of higher degree

Theorem. Let  $X/\mathbb{P}_{\mathbb{R}}^1$  be a special quadric fibration with affine equation

$$x^2 + y^2 + z^2 = u \cdot p(u).$$

Let  $p(u) = u^{2n} + \sum_{i=0}^{n-1} a_{2i} u^{2i}$ . If  $a_0 > 0$  and  $a_{2i} \geq 0$  for all  $0 < i < n$ , then  $X$  is universally  $CH_0$ -trivial.

Theorem. Let  $X/\mathbb{P}_{\mathbb{R}}^1$  be a special quadric fibration with affine equation

$$x^2 + y^2 + z^2 = u \cdot p(u).$$

Let  $p(u) = u^{2n} + \sum_{i=0}^{2n-1} a_i u^{2i}$ . There exists a nonempty open set  $U \subset \mathbb{A}^{2n}(\mathbb{R})$  such that for any  $(a_0, \dots, a_{2n-1}) \in U$ , the associated variety  $X$  is universally  $CH_0$ -trivial.

**Universal Chow triviality for  $\deg(p) = 2$  via the criterion :**

**(2b) For  $F = \mathbb{R}(\Delta)$ , the map  $H^3(F, \mathbb{Z}/2) \rightarrow H_{nr}^3(F(X)/F, \mathbb{Z}/2)$  is an isomorphism.**

*Theorem. Let  $p(u) \in \mathbb{R}[u]$  be a positive polynomial of degree 2. Let  $X/\mathbb{P}_{\mathbb{R}}^1$  be a special quadric fibration with affine equation*

$$x^2 + y^2 + z^2 = u.p(u).$$

*Assume that the elliptic curve  $E/\mathbb{R}$  defined by  $z^2 = u.p(u)$  has “odd” complex multiplication, namely  $\text{End}_{\mathbb{C}} E = \mathbb{Z}[\omega]$ , with  $\omega^2 - d\omega + c = 0$ ,  $c, d \in \mathbb{Z}$  and  $d$  odd. Let  $\Delta$  be defined by  $w^2 = v.p(-v)$ . Let  $F = \mathbb{R}(\Delta)$ . Then the map  $H^3(F, \mathbb{Z}/2) \rightarrow H_{nr}^3(F(X)/F, \mathbb{Z}/2)$  is an isomorphism, and the variety  $X$  is universally  $CH_0$ -trivial.*

### *Main points of the proof*

Let  $F$  be any overfield of  $\mathbb{R}$ . We consider the birational conic bundle fibration  $X_F \rightarrow \mathbb{P}_F^2$  induced by the projection map

$$(x, y, z, u) \mapsto (z, u) \in \mathbb{A}^2 \subset \mathbb{P}^2.$$

The fibration is ramified along the elliptic curve  $E_F \subset \mathbb{P}_F^2$  with affine equation  $z^2 = u \cdot p(u)$  and possibly along the line at infinity. By general  $K$ -theory results on conics, and a standard analysis of residues and their functoriality, one shows that any class  $\beta \in H_{nr}^3(F(X)/F, \mathbb{Z}/2)$  trivial at an  $F$ -point is the image of a class  $\alpha \in H^3(F(\mathbb{P}^2), \mathbb{Z}/2)$  whose residues away from  $E_F$  and the line at infinity of  $\mathbb{P}_F^2$  are zero, and whose residue at  $E_F$  belongs to  $\text{Ker}[\text{Br}(E_F) \rightarrow \text{Br}(E_{F'})]$ , where  $F' := F(\sqrt{-1})$ . *We would like to get rid of this possible residue.*

Let  $G = \mathbb{Z}/2 = \text{Gal}(F'/F)$ . We have a standard exact sequence

$$0 \rightarrow H^2(G, F') \rightarrow \text{Ker}[\text{Br}(E_F) \rightarrow \text{Br}(E_{F'})] \rightarrow H^1(G, \text{Pic}(E_{F'})) \rightarrow 0.$$

Let  $F = \mathbb{R}(\Delta)$ . The  $G$ -lattice  $M := \text{Hom}_{\mathbb{C}}(\Delta_{\mathbb{C}}, E_{\mathbb{C}})$  is isomorphic, as an abelian group, to  $\text{End}_{\mathbb{C}}(E_{\mathbb{C}})$ , which is abstractly  $\mathbb{Z}$  or  $\mathbb{Z} \oplus \mathbb{Z}$  (last case, complex multiplication).

Using  $E(\mathbb{R})$  connected, one gets  $H^1(G, \text{Pic}(E_{F'})) \simeq H^1(G, M)$ .

So the obstruction to solving our problem lies in the finite group  $H^1(G, M)$ , which one may show is either 0 or  $\mathbb{Z}/2$ .

Key technical result :

Proposition. Assume that the elliptic curve  $E/\mathbb{R}$  defined by  $z^2 = u.p(u)$  has “odd” complex multiplication, namely  $\text{End}_{\mathbb{C}} E = \mathbb{Z}[\omega]$ , with  $\omega^2 - d\omega + c = 0$ ,  $c, d \in \mathbb{Z}$  and  $d$  odd. Let  $\Delta$  be defined by  $w^2 = v.p(-v)$ . Let  $F = \mathbb{R}(\Delta)$  and  $F' = F(\sqrt{-1})$ . Then  $H^1(G, \text{Pic}(E_{F'})) = H^1(G, M) = 0$ .

Under this hypothesis, the residue of  $\alpha$  at  $E_F$  is of the shape  $(\delta, -1)$  with  $\delta \in F^\times$ . Over  $\mathbb{A}_F^2$  the classes  $\alpha$  and  $(\delta, z^2 - up(u), -1)$  have the same residues. Their difference is thus in  $H^3(F, \mathbb{Z}/2)$ . Since  $-(z^2 - up(u)) = x^2 + y^2$  in  $\mathbb{R}(X)$ , the image of  $(\delta, z^2 - up(u), -1)$  in  $H^3(F(X), \mathbb{Z}/2)$  is  $(\delta, -1, -1)$  hence comes from  $H^3(F, \mathbb{Z}/2)$ .

QED

## Comparing the two methods for $\deg(p) = 2$

Let  $E/\mathbb{R}$  be the elliptic curve with equation  $z^2 = u.(u^2 + au + b)$ ,  $a, b \in \mathbb{R}$ . We assume  $b > 0$  and  $0 \leq a^2/b < 4$ .

One computes

$$j(E) = 256[3 - (a^2/b)]^3/[4 - (a^2/b)] \in \mathbb{R}.$$

$0 \leq a^2/b \leq 3$  if and only if  $j(E) \geq 0$ , and then  $0 \leq j(E) \leq 1728$ .

$3 \leq a^2/b < 4$  if and only if  $j(E) \leq 0$ .

$a^2/b = 3$  corresponds to  $j(E) = 0$  and  $a^2/b = 0$  to  $j(E) = 1728$



Chow triviality for  $x^2 + y^2 + z^2 = u \cdot (u^2 + au + b)$ , with  $b > 0$  and  $a^2 - 4b < 0$ .

First method (sum of squares)

This works for all  $(a, b)$ , with  $b > 0$ ,  $0 \leq a^2/b \leq 3$ , i.e. any  $j(E) \geq 0$ . Here  $j(E)$  takes all values in  $[0, 1728]$ .

Second method (conic bundle fibration and  $E$  had odd complex multiplication). For  $3 < a^2/b < 4$ , i.e.  $j(E) < 0$ , this is the only method we have. So we do not know what happens if such  $E$  does not have complex multiplication or has even complex multiplication.

For  $E$  with odd complex multiplication, the invariant  $j(E) \in \mathbb{R}$  is algebraic, the values it takes are in  $[-\infty, 1728]$ .

Theorem (Yu. Zarhin) : *these values are dense in  $[-\infty, 1728]$ .*

In a recent paper, Zarhin systematically analyzes odd versus even complex multiplication.

## Examples for which we can prove $X$ is universally $CH_0$ -trivial

$$\rho(u) = u^2 - 3u + 3$$

$E$  is given by  $z^2 = (u - 1)^3 + 1$ . It has complex multiplication by  $\omega$  with  $\omega^2 + \omega + 1 = 0$ . This it has odd CM. Here  $j(E) = 0$ . Both methods apply.

$$\rho(u) = u^2 + 1$$

$E$  is given by  $z^2 = u(u^2 + 1)$ . It has  $j(E) = 1728$ . The first method applies. The curve  $E$  has CM by  $\omega = \sqrt{-1}$ , but  $\omega^2 + 1 = 0$  hence it is not odd CM. The second method does not apply.

$\rho(u) = u^2 - 21u + 112$ . Here  $j(E) < 0$ , the first method does not apply. The curve has complex multiplication by  $\mathbb{Z}[\omega]$  with  $\omega = (1 + \sqrt{-7})/2$ . Here  $\omega^2 - \omega + 2 = 0$ , thus is odd CM, the second method applies.

## Open problems

Let  $X/\mathbb{R}$  be a smooth projective model of the variety with affine equation  $x^2 + y^2 + z^2 = u.p(u)$ , with  $p(u)$  monic, separable, positive on  $\mathbb{R}$ , of degree at least 2. Let  $\Delta/\mathbb{R}$  be the curve with affine equation  $w^2 = v.p(-v)$ .

### Are the following equivalent conditions always satisfied?

- The variety  $X/\mathbb{R}$  is universally  $CH_0$ -trivial.
- The rational function  $u + v \in \mathbb{R}(X \times_{\mathbb{R}} \Delta)$  (a sum of 6 squares) is a sum of 4 squares.

- Are there examples for which  $X$  is rational over  $\mathbb{R}$ ?
- Are there examples for which  $X$  is not rational over  $\mathbb{R}$ ?
- What about  $\deg(p) = 2$ ?
- What about  $x^2 + y^2 + z^2 = u.(u^2 + 1)$ ? (universally  $CH_0$ -trivial).